



University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Nigel Stallard

Article Title: An adaptive seamless phase II/III clinical trial design incorporating short-term endpoint information

Year of publication: 2012

Link to published article:

<http://dx.doi.org/10.1186/1745-6215-12-S1-A2>

Publisher statement: None

ORAL PRESENTATION

Open Access

An adaptive seamless phase II/III clinical trial design incorporating short-term endpoint information

Nigel Stallard

From Clinical Trials Methodology Conference 2011
Bristol, UK. 4-5 October 2011

Adaptive seamless phase II/III designs enable a clinical trial to be conducted in stages with the most promising of a number of experimental treatments selected on the basis of data observed in the first stage to continue along with the control treatment to the second and any subsequent stages. The main statistical challenge in such a design is ensuring control of the type I error rate. Most methodology for such trials is based on the same endpoint being used for interim and final analyses. In some settings the primary endpoint can be observed only after long-term follow-up. In this case it may be desirable to use short-term endpoint data along with any long-term data available at the interim analysis to inform treatment selection. If short-term data are available for some patients for whom the primary endpoint is not available, basing treatment selection on these data may lead to inflation of the type I error rate.

This talk presents a new method [1] that allows the use of short-term endpoint information for treatment selection whilst controlling the overall type I error rate. The method builds on the work of Galbraith and Marschner [2], who proposed a method for using short-term endpoints in interim analyses for comparison of a single experimental treatment with a control. The method is based on adjustment of the usual group-sequential boundaries [3,4] to allow for monitoring of a test statistic obtained from fitting a bivariate normal model to the correlated short-term and long-term endpoints. The model can be fitted either using a linear mixed model or by combining the results from separate

linear regression models for the short and long-term data [5].

Result will be presented from a simulation study to investigate the properties of the new method. In addition to verifying control of the type I error rate, these results show that the use of the short-term endpoint data can lead to an increase in power when the short and long-term endpoints are correlated.

Published: 13 December 2011

References

1. Stallard N: A confirmatory seamless phase II/III clinical trial design incorporating short-term endpoint information. *Statistics in Medicine* 2010, **29**:959-971.
2. Galbraith S, Marschner IC: Interim analysis of continuous long-term endpoints in clinical trials with longitudinal outcomes. *Statistics in Medicine* 2003, **22**:1787-1805.
3. Jennison C, Turnbull BW: Group-sequential analysis incorporating covariate information. *Journal of the American Statistical Association* 1997, **92**:1330-1341.
4. Stallard N, Todd S: Sequential designs for phase III clinical trials incorporating treatment selection. *Statistics in Medicine* 2003, **22**:689-703.
5. Engel B, Walstra P: Increasing precision or reducing expense in regression experiments by using information from a concomitant variable. *Biometrics* 1991, **47**:13-20.

doi:10.1186/1745-6215-12-S1-A2

Cite this article as: Stallard: An adaptive seamless phase II/III clinical trial design incorporating short-term endpoint information. *Trials* 2011 **12** (Suppl 1):A2.

Correspondence: n.stallard@warwick.ac.uk
Statistics and Epidemiology, Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, CV4 7AL, UK